Rapid Detect 505

Collect sufficient sample the first time with the first ever sponge saturation indicator!

- Patent pending
- Single drug per strip
- Exclusive tests not available in any other device
- Detects up to 10 drugs simultaneously
- Tamper evident and air tight to retain remaining saliva for laboratory confirmation
- Magnification windows for clear results
- MADE IN THE USA

Oral Fluid Collector (Sponge)

Studies have proven:

- Highest rate of drug recovery without buffer
- Highest, most consistent rate of absorption
- Reduces viscosity of saliva for faster, more consistent results


Sponge Saturation Indicator

Typical time-determined collection methods are often ineffective when donors are smokers or users of certain prescription drugs, which can cause severe dry mouth. The sponge saturation indicator activates when sufficient sample has been collected.

- Reduce the cost of wasted devices due to insufficient sample common with other oral fluid tests
- Collectors invest only the time necessary to gather the sample and not a minute more



Results interpretation



Tests exclusive to this device


Cotinine: 50 ng/mlEDDP: 20 ng/mlKetamine: 50 ng/ml

Market-leading drug cutoffs

Methadone: 30 ng/mlBuprenorphine: 5 ng/ml

SALIVA CUTOFFS	
Alcohol (ALC)	.02%
Amphetamine (AMP)	50
Barbiturate (BAR)	50
Benzodiazepines (BZO)	10
Buprenorphine (BUP)	5
Cocaine (COC)	20
Cotinine (COT)	50
EDDP	20






Manufactured for:



SPONGE SATURATION INDICATOR

EXAMPLE: Presumptive positive for cocaine (COC)





FORENSIC USE OR EXPORT ONLY

301 Turman Street, Poteau, OK 74953 888-404-0020 | Fax 918-649-1005 www.rapiddetect.com